

Product Description

Pioneering GTPase and Oncogene Product Development since 2010


PDE4A RABBIT PAB

货号: S220790 产品全名: PDE4A 兔多抗 基因符号 PDE4; DPDE2; PDE46 UNIPROT ID: P27815 (Gene Accession - NP_001104777) 背景: The protein encoded by this gene belongs to the cyclic nucleotide phosphodiesterase (PDE) family, and PDE4 subfamily. This PDE hydrolyzes the second messenger, cAMP, which is a regulator and mediator of a number of cellular responses to extracellular signals. Thus, by regulating the cellular concentration of cAMP, this protein plays a key role in many important physiological processes. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. 抗原: Synthetic peptide of human PDE4A 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Signal Transduction 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 220790(PDE4A Antibody) at a dilution of 1/40(Cytoplasm).




The image on the left is immunohistochemistry of paraffinembedded Human brain tissue using 220790(Anti-PDE4A Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the synthetic peptide and then with 220790(Anti-PDE4A Antibody) at dilution 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human brain tissue is first treated with synthetic peptide and then with D262029(Anti-PDE4A Antibody) at dilution 1/40.



Lane 1-2: Hela cells, A172 cells; Primary antibody: 220790(PDE4A Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 20 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010