

## PPAR DELTA (4F4) MOUSE MAB

货号: N261392

产品全名: PPAR delta (4F4) 小鼠单抗

基因符号 FAAR; NR1C2; NUC1; Peroxisome proliferative activated receptor delta

**UNIPROT ID:** Q03181

**背景:** Ligand-activated transcription factor. Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma-linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand.

**抗原:** Purified recombinant protein expressed in E.coli.

**经过测试的应用:** IHC-P

**推荐稀释比:** IHC: 1/50-1/100

**种属反应性:** Mouse

**克隆性:** Mouse Monoclonal

**克隆编号:** 4F4-3B6-7D1

**分子量:** -

**亚型:** IgG1


**纯化:** Affinity Purified

**种属反应性:** Human,Rat,Mouse


**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**研究领域:** Epigenetics and Nuclear Signaling

**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemical analysis of paraffin-embedded Human tonsils using PPAR delta (4F4) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded Human Brain Tissue using PPAR delta (4F4) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.