

## PPAR GAMMA RABBIT PAB

货号: N225397

产品全名: PPAR gamma 兔多抗

基因符号 CIMT1; GLM1; NR1C3; Nuclear receptor subfamily 1 group C member 3; Peroxisome proliferator activated nuclear receptor gamma variant 1; Peroxisome proliferator activated receptor gamma 1; Peroxisome Proliferator Activated Receptor gamma; Peroxisome proliferator-activated receptor gamma; PPAR gamma; PPAR-gamma; PPARG; PPARG\_HUMAN; PPARG1; PPARG2; PPARgamma.

**UNIPROT ID:** P37231

**背景:** Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis.

**抗原:** The antiserum was produced against synthesized peptide derived from human PPAR-gamma. AA range:78-127

**经过测试的应用:** ICC/IF,WB,IHC-F,IHC-P,ELISA

**推荐稀释比:** WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ELISA: 1/10000

**种属反应性:** Rabbit

**克隆性:** Rabbit Polyclonal

**分子量:** Calculated MW: 58 kDa; Observed MW: 58 kDa

**亚型:** IgG


**纯化:** Affinity Purified

**种属反应性:** Human, Mouse and Rat


**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**研究领域:** Epigenetics and Nuclear Signaling

**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded rat spleen tissue using PPAR gamma antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Sample with blocking peptide on the right.



Western blot analysis of PPAR gamma in HuvEc lysates using PPAR gamma antibody.