

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

PRCP RABBIT PAB

货号: S216743

产品全名: PRCP 兔多抗 基因符号 PCP; HUMPCP

UNIPROT ID: P42785 (Gene Accession - BC001500)

背景: The protein encoded by this gene is a lysosomal prolylcarboxypeptidase, which cleaves C-terminal amino acids linked to proline in peptides such as angiotension II, III and des-Arg9-bradykinin. The cleavage occurs at acidic pH, but the enzyme activity is retained with some substrates at neutral pH. This enzyme has been shown to be an activator of the cell matrix-associated prekallikrein. The importance of angiotension II, one of the substrates of this enzyme, in regulating blood pressure and electrolyte balance suggests that this gene may be related to essential hypertension. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

抗原: Fusion protein of human PRCP 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cell Biology, Cardiovascular

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 216743(PRCP Antibody) at a dilution of 1/15(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 216743(Anti-PRCP Antibody) at dilution 1/15.



The image on the left is immunohistochemistry of paraffinembedded Human ovarian cancer tissue using 216743(Anti-PRCP Antibody) at a dilution of 1/15.



In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with fusion protein and then with D221156(Anti-PRCP Antibody) at dilution 1/15.



Gel: 10%SDS-PAGE, Lysate: 50 µg; Lane 1-3: SKOV3 cells, A549 cells, human liver cancer tissue; Primary antibody: 216743(PRCP Antibody) at dilution 1/100; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010