

PSMB7 RABBIT PAB

货号: S217094

产品全名: PSMB7 兔多抗

基因符号 Z

UNIPROT ID: Q99436 (Gene Accession - BC000509)

背景: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. The encoded protein is a member of the proteasome B-type family, also known as the T1B family, and is a 20S core beta subunit in the proteasome. Expression of this catalytic subunit is downregulated by gamma interferon, and proteolytic processing is required to generate a mature subunit. A pseudogene of this gene is located on the long arm of chromosome 14.

抗原: Fusion protein of human PSMB7

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse, Rat


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cell Biology


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human colon cancer tissue using 217094(PSMB7 Antibody) at a dilution of 1/25(Cytoplasm and Nucleus).




In comparison with the IHC on the left, the same paraffin-embedded Human colon cancer tissue is first treated with the fusion protein and then with 217094(Anti-PSMB7 Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using 217094(Anti-PSMB7 Antibody) at a dilution of 1/25.



In comparison with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with fusion protein and then with D221776(Anti-PSMB7 Antibody) at dilution 1/25.



Gel: 10%SDS-PAGE, Lysate: 40 μ g;
Lane 1-4: 293T cells, hepG2 cells, human testis tissue, Human lung cancer tissue;
Primary antibody: 217094(PSMB7 Antibody) at dilution 1/310;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 5 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
