

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

RBBP8 RABBIT PAB

货号: S217312

产品全名: RBBP8 兔多抗

基因符号 RIM; COM1; CTIP; JWDS; SAE2; SCKL2

UNIPROT ID: Q99708 (Gene Accession - BC030590)

背景: The protein encoded by this gene is a ubiquitously expressed nuclear protein. It is found among several proteins that bind directly to retinoblastoma protein, which regulates cell proliferation. This protein complexes with transcriptional co-repressor CTBP. It is also associated with BRCA1 and is thought to modulate the functions of BRCA1 in transcriptional regulation, DNA repair, and/or cell cycle checkpoint control. It is suggested that this gene may itself be a tumor suppressor acting in the same pathway as BRCA1.

抗原: Fusion protein of human RBBP8

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling, Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 217312(RBBP8 Antibody) at a dilution of 1/45(cytoplasm or nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 217312(Anti-RBBP8 Antibody) at dilution 1/45.



The image on the left is immunohistochemistry of paraffinembedded Human breast cancer tissue using 217312(Anti-RBBP8 Antibody) at a dilution of 1/45.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with fusion protein and then with D222163(Anti-RBBP8 Antibody) at dilution 1/45.



Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane 1-3: Hepg2, Hela and 231 cell; Primary antibody: 217312(RBBP8 Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 2 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010