

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010


## **REL RABBIT PAB**

货号: S217305 产品全名: REL 兔多抗 基因符号 C-Rel UNIPROT ID: Q04864 (Gene Accession - NP\_002899) 背景: The REL gene encodes c-Rel, a transcription factor that is a member of the Rel/NFKB family, which also includes RELA (MIM 164014), RELB (604758), NFKB1 (MIM 164011), and NFKB2 (MIM 164012). These proteins are related through a highly conserved N-terminal region termed the 'Rel domain, which is responsible for DNA binding, dimerization, nuclear localization, and binding to the NFKB inhibitor 抗原: Fusion protein of human REL 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 100-300;WB: 1000-5000;ELISA: 2000-10000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Signal Transduction, Epigenetics and Nuclear Signaling, Cancer 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 217305(REL Antibody) at a dilution of 1/50(Nucleus or Cytoplasm).



The image on the left is immunohistochemistry of paraffinembedded Human brain tissue using 217305 (Anti-REL Antibody) at a Human brain tissue is first treated with fusion protein and then with dilution of 1/50.



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the fusion protein and then with 217305(Anti-REL Antibody) at dilution 1/50.



In comparision with the IHC on the left, the same paraffin-embedded D222147(Anti-REL Antibody) at dilution 1/50.



Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Hela cells, 293T cells; Primary antibody: 217305(REL Antibody) at dilution 1/1050; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010