

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

RFWD3 RABBIT PAB

货号: N225401

产品全名: RFWD3 兔多抗 基因符号 FANCW; RNF201 UNIPROTID: O6PCD5

背景: E3 ubiquitin-protein ligase required for the repair of DNA interstrand cross-links (ICL) in response to DNA damage (PubMed:21504906, PubMed:21558276, PubMed:26474068, PubMed:28575657,

PubMed:28575658). Plays a key role in RPA-mediated DNA damage signaling and repair (PubMed:21504906, PubMed:21558276, PubMed:26474068, PubMed:28575657, PubMed:28575658,

PubMed:28691929). Acts by mediating ubiquitination of the RPA complex (RPA1, RPA2 and RPA3 subunits) and RAD51 at stalled replication forks, leading to remove them from DNA damage sites and promote homologous recombination (PubMed:26474068, PubMed:28575657, PubMed:28575658). Also mediates the ubiquitination of p53/TP53 in the late response to DNA damage, and acts as a positive regulator of p53/TP53 stability, thereby regulating the G1/S DNA damage checkpoint (PubMed:20173098). May act by catalyzing the formation of short polyubiquitin chains on p53/TP53 that are not targeted to the proteasome (PubMed:20173098). In response to ionizing radiation, interacts with MDM2 and enhances p53/TP53 ubiquitination, possibly by restricting MDM2 from extending polyubiquitin

抗原: The antiserum was produced against synthesized peptide derived from human RFWD3. AA range:374-423

经过测试的应用: WB,IHC-P,ELISA

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000

chains on ubiquitinated p53/TP53 (PubMed:20173098).

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 85 kDa; Observed MW: 85 kDa

亚型: IgG

纯化: Affinity Chromatography

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.3 containing 50% glycerol,

0.5% BSA and 0.02% sodium azide


研究领域: -

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Western blot analysis of RFWD3 in HepG2 lysates using RFWD3 antibody.



Western blot analysis of RFWD3 in 823 lysates using RFWD3 antibody.



Western blot analysis of RFWD3 in MCF-7 lysates using RFWD3 antibody. The lane on the right is blocked with the synthesized peptide.