

SCRN2 RABBIT PAB

货号: S218879

产品全名: SCRN2 兔多抗

基因符号: Ses2

UNIPROT ID: Q96FV2 (Gene Accession - BC017317)

背景: SCRN genes encode protein members of the secernin family including Secernin 1, Secernin 2 and Secernin 3. Function of Secernin 2 and Secernin 3 are so far not well understood. Secernin 1 is a novel 50-kDa cytosolic protein that appears to be involved in the regulation of exocytosis from peritoneal mast cells. Secernin-1 a novel tumor-associated antigen (TAA) and may be a universal marker of different cancer types including gastric cancer, which has been validated by several researches. Measurement of SCRNI can be used in the early detection of cancer or in the surveillance of patients who undergo surgery.

抗原: Fusion protein of human SCRN2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse, Rat


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Cell Biology


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human lung cancer tissue using 218879(SCR2N2 Antibody) at a dilution of 1/60(Nucleus or Cytoplasm).



In comparison with the IHC on the left, the same paraffin-embedded Human lung cancer tissue is first treated with the fusion protein and then with 218879(Anti-SCR2N2 Antibody) at dilution 1/60.




The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using 218879(Anti-SCR2N2 Antibody) at a dilution of 1/60.



In comparison with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with fusion protein and then with D225426(Anti-SCR2N2 Antibody) at dilution 1/60.

kDa



Gel: 8%SDS-PAGE, Lysate: 40 µg;
 Lane: Mouse small intestines tissue lysate;
 Primary antibody: 218879(SCR2N2 Antibody) at dilution 1/400;
 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
 Exposure time: 10 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
