

Product Description

Pioneering GTPase and Oncogene Product Development since 2010


SEC14L2 RABBIT PAB

货号: S220888 产品全名: SEC14L2 兔多抗 基因符号 SPF; TAP; TAP1; C22orf6 UNIPROT ID: O76054 (Gene Accession - NP_001191133) 背景: This gene encodes a cytosolic protein which belongs to a family of lipid-binding proteins including Sec14p, alpha-tocopherol transfer protein, and cellular retinol-binding protein. The encoded protein stimulates squalene monooxygenase which is a downstream enzyme in the cholesterol biosynthetic pathway. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. 抗原: Synthetic peptide of human SEC14L2 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Metabolism 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 220888(SEC14L2 Antibody) at a dilution of 1/40(Cytoplasm).




The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 220888(Anti-SEC14L2 Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 220888(Anti-SEC14L2 Antibody) at dilution 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with synthetic peptide and then with D262176(Anti-SEC14L2 Antibody) at dilution 1/40.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-3: Mouse liver tissue, PC3 cells, human fetal liver tissue; Primary antibody: 220888(SEC14L2 Antibody) at dilution 1/400; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010