

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

SLC1A4 RABBIT PAB

货号: S219990

产品全名: SLC1A4 兔多抗 基因符号 SATT, ASCT1

UNIPROT ID: P43007 (Gene Accession - NP_003029)

背景: Neutral amino acid transporter A is a protein that in humans is encoded by the SLC1A4 gene. In melanocytic cells SLC1A4 gene expression may be regulated by MITF. ASCTI does not transport glutamate or aspartate but alanine, serine, cysteine, and threonine instead. ASCTI transporter functions primarily as an amino acid exchanger. Transport is associated with a chloride channel activity that is thermodynamically uncoupled from amino acid transport.

抗原: Synthetic peptide of human SLC1A4

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 1000-5000;ELISA: 2000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 219990(SLC1A4 Antibody) at a dilution of 1/70(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with the synthetic peptide and then with 219990(Anti-SLC1A4 Antibody) at dilution 1/70.



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 219990(Anti-SLC1A4 Antibody) at a dilution of 1/70.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with synthetic peptide and then with D260739(Anti-SLC1A4 Antibody) at dilution 1/70.



Gel: 10+12%SDS-PAGE, Lysate: 30 µg; Lane: Human liver cancer tissue; Primary antibody: 219990(SLC1A4 Ant

Primary antibody: 219990(SLC1A4 Antibody) at dilution 1/1250; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 20 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010