

## SMAD5 RABBIT MAB

货号: N262912

产品全名: Smad5 兔单克隆抗体

基因符号 DKFZp781C1895; DKFZp781O1323; Dwfc; hSmad 5; hSmad5; JV5 1; JV5-1; MAD homolog 5; MAD mothers against decapentaplegic homolog 5; MAD; mothers against decapentaplegic homolog 5; MADH 5; MADH5; Mothers against decapentaplegic homolog 5; Mothers against DPP homolog 5; MusMLP; SMA and MAD related protein 5; SMAD 5; SMAD family member 5; SMAD mothers against DPP homolog 5; Smad5; Smad5; SMAD5\_HUMAN.

**UNIPROT ID:** Q99717

**背景:** Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. Smad5 is a receptor-regulated Smad (R-Smad). Smad5 is required for normal development of the cardiovascular system in vivo; lack of the Smad5 gene results in apoptosis of cardiac myocytes. 3 Upregulation of Smad5 has been reported to mediate apoptosis of gastric epithelial cells induced by Helicobacter pylori infection. Tissue specificity: Ubiquitous.

**抗原:** A synthetic peptide of human SMAD5

**经过测试的应用:** WB,IHC-F,IHC-P,ICC/IF

**推荐稀释比:** WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200

**种属反应性:** Rabbit

**克隆性:** Rabbit Monoclonal

**克隆编号:** R07-2A5

**分子量:** Calculated MW: 52 kDa; Observed MW: 60 kDa

**亚型:** IgG


**纯化:** Affinity Purified

**种属反应性:** Human,Rat


**成分:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.3 containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

**研究领域:** Signal Transduction

**储存和运输:** Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using SMAD5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of Smad5 in C6, HeLa lysates using Smad5 antibody.