

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

SNIP1 RABBIT PAB

货号: S219380 产品全名: SNIP1 兔多抗 基因符号 PML1; PMRED

UNIPROT ID: Q8TAD8 (Gene Accession - BC027040)

背景: This gene encodes a protein that contains a coiled-coil motif and C-terminal forkhead-associated (FHA) domain. The encoded protein functions as a transcriptional coactivator that increases c-Myc activity and inhibits transforming growth factor beta (TGF-beta) and nuclear factor kappa-B (NF-kB) signaling. The encoded protein also regulates the stability of cyclin DI mRNA, and may play a role in cell proliferation and cancer progression. Mutations in this gene are a cause of psychomotor retardation, epilepsy, and craniofacial dysmorphism (PMRED).

抗原: Fusion protein of human SNIP1 经过测试的应用: ELISA, WB, IHC


推荐稀释比: IHC: 50-200;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal


亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 219380(SNIP1 Antibody) at a dilution of 1/80(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 219380(Anti-SNIP1 Antibody) at dilution 1/80.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane: TM4 cell lysate;

Primary antibody: 219380(SNIP1 Antibody) at dilution 1/1150;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000

dilution;

Exposure time: 30 seconds