

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

SP110 RABBIT PAB

货号: S219499

产品全名: SP110 兔多抗

基因符号 IPR1; VODI; IFI41; IFI75

UNIPROT ID: Q9HB58 (Gene Accession - BC019059)

背景: The nuclear body is a multiprotein complex that may have a role in the regulation of gene transcription. This gene is a member of the SP100/SP140 family of nuclear body proteins and encodes a leukocyte-specific nuclear body component. The protein can function as an activator of gene transcription and may serve as a nuclear hormone receptor coactivator. In addition, it has been suggested that the protein may play a role in ribosome biogenesis and in the induction of myeloid cell differentiation. Alternative splicing has been observed for this gene and three transcript variants, encoding distinct isoforms, have been identified.

抗原: Fusion protein of human SP110 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-100;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human gastric cancer tissue using 219499(SP110 Antibody) at a dilution of 1/135(Nucleus).


In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with the fusion protein and then with 219499(Anti-SP110 Antibody) at dilution 1/135.



The image on the left is immunohistochemistry of paraffinembedded Human tonsil tissue using 219499(Anti-SP110 Antibody) at a dilution of 1/135.



In comparision with the IHC on the left, the same paraffin-embedded Human tonsil tissue is first treated with fusion protein and then with D227390(Anti-SP110 Antibody) at dilution 1/135.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: LOVO cell lysate; Primary antibody: 219499(SP110 Antibody) at dilution 1/1000; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 30 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010