

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ST8SIA4 RABBIT PAB

货号: S218983

产品全名: ST8SIA4 兔多抗

基因符号 PST; PST1; SIAT8D; ST8SIA-IV

UNIPROT ID: Q92187 (Gene Accession - BC027866)

背景: The protein encoded by this gene catalyzes the polycondensation of alpha-2,8-linked sialic acid required for the synthesis of polysialic acid, a modulator of the adhesive properties of neural cell adhesion molecule (NCAMI). The encoded protein, which is a member of glycosyltransferase family 29, is a type II membrane protein that may be present in the Golgi apparatus. Two transcript variants encoding different isoforms have been found for this gene.

抗原: Fusion protein of human ST8SIA4

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cell Biology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 218983(ST8SIA4 Antibody) at a dilution of 1/50(Cytoplasm and Nucleus).


In comparision with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with the fusion protein and then with 218983(Anti-ST8SIA4 Antibody) at dilution 1/50.




In comparision with the IHC on the left, the same paraffin-embedded



The image on the left is immunohistochemistry of paraffinembedded Human breast cancer tissue using 218983(Anti-ST8SIA4 Antibody) at a dilution of 1/50.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with fusion protein and then with D225604(Anti-ST8SIA4 Antibody) at dilution 1/50.



kDa

Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Human cerebella tissue lysate; Primary antibody: 218983(ST8SIA4 Antibody) at dilution 1/350; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 50 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010