

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **STAT5A RABBIT PAB**

货号: S216260

产品全名: STAT5A 兔多抗 基因符号 MGF; STAT5

UNIPROT ID: P42229 (Gene Accession - BC027036)

背景: The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCl2l1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have been found for this gene.

抗原: Fusion protein of human STAT5A

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 500-5000


种属反应性: Rabbit

克隆性: Rabbit Polyclonal


亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction, Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using 216260(STAT5A Antibody) at a dilution of 1/25(Cytoplasm).



Immunohistochemistry analysis of paraffin-embedded Human ovarian cancer tissue using 216260(Anti-STAT5A Antibody) at a dilution of 1/25



Gel: 10%SDS-PAGE, Lysate: 30 µg; Lane: Human fetal lung tissue lysate;

Primary antibody: 216260(STAT5A Antibody) at dilution 1/450; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 30 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010