

STK32B RABBIT PAB

货号: S219385

产品全名: STK32B 兔多抗

基因符号 STK32; STKG6; YANK2; HSA250839

UNIPROT ID: Q9NY57 (Gene Accession - BC038238)

背景: This gene encodes a serine-threonine protein kinase. Serine-threonine kinases transfer phosphate molecules to the oxygen atoms of serine and threonine. A genomic deletion affecting this gene has been associated with Ellis-van Creveld syndrome, an autosomal recessive skeletal dysplasia. Alternative splicing results in multiple transcript variants.

抗原: Fusion protein of human STK32B

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 100-300;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse

成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 219385(STK32B Antibody) at a dilution of 1/160(Cytoplasm and Nucleus).

In comparison with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 219385(Anti-STK32B Antibody) at dilution 1/160.



Gel: 8%SDS-PAGE, Lysate: 40 µg;

Lane 1-2: Mouse liver tissue and Mouse brain tissue lysates;

Primary antibody: 219385(STK32B Antibody) at dilution 1/1000;

Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 3 minutes