

STT3A RABBIT PAB

货号: S218774

产品全名: STT3A 兔多抗

基因符号 TMC; ITMI; STT3-A

UNIPROT ID: P46977 (Gene Accession - BC020965)

背景: The protein encoded by this gene is a catalytic subunit of the N-oligosaccharyltransferase (OST) complex, which functions in the endoplasmic reticulum to transfer glycan chains to asparagine residues of target proteins. A separate complex containing a similar catalytic subunit with an overlapping function also exists. Multiple transcript variants encoding different isoforms have been found for this gene.

抗原: Fusion protein of human STT3A

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 30-150; WB: 200-1000; ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse

成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cell Biology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 218774 (STT3A Antibody) at a dilution of 1/30 (Cytoplasm).

In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 218774 (Anti-STT3A Antibody) at dilution 1/30.



Gel: 8% SDS-PAGE, Lysate: 40 µg;

Lane 1-2: Mouse liver tissue and Human placenta tissue lysates;

Primary antibody: 218774 (STT3A Antibody) at dilution 1/200;

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 10 seconds