

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **SYCE1 RABBIT PAB**

货号: S219562

产品全名: SYCEI 兔多抗

基因符号 CT76; POF12; SPGF15; C10orf94

UNIPROT ID: Q8N0S2 (Gene Accession - BC034821)

背景: This gene encodes a member of the synaptonemal comple,x which links homologous chromosomes during prophase I of meiosis. The tripartite structure of the complex is highly conserved amongst metazoans. It consists of two lateral elements and a central region formed by transverse elements and a central element. The protein encoded by this gene localizes to the central element and is required for initiation and elongation of the synapsis. Allelic variants of this gene have been associated with premature ovarian failure and spermatogenic failure. Alternative splicing results in multiple transcript variants.

抗原: Fusion protein of human SYCEI

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cell Biology, Epigenetics and Nuclear Signaling

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human colorectal cancer tissue using 219562(SYCEI Antibody) at a dilution of Human colorectal cancer tissue is first treated with the fusion protein 1/55(Cytoplasm).


In comparision with the IHC on the left, the same paraffin-embedded



The image on the left is immunohistochemistry of paraffinembedded Human liver cancer tissue using 219562(Anti-SYCE) Antibody) at a dilution of 1/55.



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with fusion protein and then with D227772(Anti-SYCE1 Antibody) at dilution 1/55.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-3: Hela, HepG2, LO2 cell lysates; Primary antibody: 219562(SYCEI Antibody) at dilution 1/800; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 5 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010