

TACR2 RABBIT PAB

货号: S220144

产品全名: TACR2 兔多抗

基因符号: SKR, NK2R, NKNAR, TAC2R

UNIPROT ID: P21452 (Gene Accession - NP_001048)

背景: This gene belongs to a family of genes that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin neuropeptide substance K, also referred to as neurokinin A.

抗原: Synthetic peptide of human TACR2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: Oct-50;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Neuroscience


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 220144(TACR2 Antibody) at a dilution of 1/10(Cytoplasm).




In comparison with the IHC on the left, the same paraffin-embedded Human ovarian cancer tissue is first treated with the synthetic peptide and then with 220144(Anti-TACR2 Antibody) at dilution 1/10.



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using 220144(Anti-TACR2 Antibody) at a dilution of 1/10.



In comparison with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with synthetic peptide and then with D261034(Anti-TACR2 Antibody) at dilution 1/10.



Gel: 10%SDS-PAGE, Lysate: 40 µg;
Lane: Human liver cancer tissue;
Primary antibody: 220144(TACR2 Antibody) at dilution 1/200;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
