

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

TAGLN2 RABBIT PAB

货号: S216786

产品全名: TAGLN2 兔多抗

基因符号 HA1756

UNIPROT ID: P37802 (Gene Accession - BC009357)

背景: The protein encoded by this gene is similar to the protein transgelin, which is one of the earliest markers of differentiated smooth muscle. The specific function of this protein has not yet been determined, although it is thought to be a tumor suppressor. Multiple transcript variants encoding different isoforms have been found for this gene.

抗原: Fusion protein of human TAGLN2

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 200-400;WB: 1000-5000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Cancer

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human ovarian cancer tissue using 216786(TAGLN2 Antibody) at a dilution of 1/240(Cytoplasm).


Human ovarian cancer tissue is first treated with the fusion protein and then with 216786(Anti-TAGLN2 Antibody) at dilution 1/240.




In comparision with the IHC on the left, the same paraffin-embedded



The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue using 216786(Anti-TAGLN2 Antibody) at a dilution of 1/240.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with fusion protein and then with D221236(Anti-TAGLN2 Antibody) at dilution 1/240.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-4: MCF7 cell, Mouse lung tissue, Hela and A375 cell lysates; Primary antibody: 216786(TAGLN2 Antibody) at dilution 1/1000; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution;

Exposure time: 2 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010