

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## TBC1D22A RABBIT PAB

货号: S217882

产品全名: TBC1D22A 兔多抗 基因符号 C22orf4; HSC79E021

UNIPROT ID: Q8WUA7 (Gene Accession - BC002743)

背景: TBC1D22A (TBC1 domain family, member 22A), also known as C22orf4, is a 517 amino acid protein that contains one Rab-GAP TBC domain and is thought to function as a GTPase-activating protein for Rab family members. Multiple isoforms of TBC1D22A exist due to alternative splicing events. The gene encoding TBC1D22A maps to human chromosome 22, which houses over 500 genes and is the second smallest human chromosome.

抗原: Fusion protein of human TBC1D22A

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 217882(TBC1D22A Antibody) at a dilution of 1/20(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the fusion protein and then with 217882(Anti-TBC1D22A Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human prostate cancer tissue using 217882(Anti-TBC1D22A Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human prostate cancer tissue is first treated with fusion protein and then with D223293(Anti-TBCID22A Antibody) at dilution 1/20.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-2: A431 cell and human lung cancer tissue; Primary antibody: 217882(TBCID22A Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 minute



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010