

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

TBP RABBIT PAB

货号: S217881 产品全名: TBP 兔多抗 基因符号 HDL4; GTF2D; SCA17; TFIID; GTF2D1 UNIPROT ID: P20226 (Gene Accession - BC109054) 背景: This gene encodes TBP, the TATA-binding prot

背景: This gene encodes TBP, the TATA-binding protein. A distinctive feature of TBP is a long string of glutamines in the N-terminus. This region of the protein modulates the DNA binding activity of the C terminus, and modulation of DNA binding affects the rate of transcription complex formation and initiation of transcription. The number of CAG repeats encoding the polyglutamine tract is usually 32-39, and expansion of the number of repeats increases the length of the polyglutamine string and is associated with spinocerebellar ataxia 17, a neurodegenerative disorder classified as a polyglutamine disease.

抗原: Full length fusion 蛋白 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG

纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 217881(TBP Antibody) at a dilution of 1/20(Nucleus).




The image on the left is immunohistochemistry of paraffinembedded Human gastric cancer tissue using 217881(Anti-TBP Antibody) at a dilution of 1/20.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-4: 231 cell, Human testis tissue, Hela and Hepg2 cell; Primary antibody: 217881(TBP Antibody) at dilution 1/400; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds

In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the fusion protein and then with 217881(Anti-TBP Antibody) at dilution 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with fusion protein and then with D223289(Anti-TBP Antibody) at dilution 1/20.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010