

Product Description

Pioneering GTPase and Oncogene Product Development since 2010


TDP2 RABBIT PAB

货号: S220534 产品全名: TDP2 兔多抗 基因符号 EAP2; AD022; EAPII; TTRAP; hTDP2; dJ30M3.3 UNIPROT ID: 095551 (Gene Accession - NP_057698) 背景: This gene encodes a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor associated factors (TRAFs), and inhibits nuclear factor-kappa-B activation. This protein has sequence and structural similarities with APEI endonuclease, which is involved in both DNA repair and the activation of transcription factors. 抗原: Synthetic peptide of human TDP2 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 50-300;WB: 500-2000;ELISA: 5000-10000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Epigenetics and Nuclear Signaling, Cancer 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human cervical cancer tissue using 220534(TDP2 Antibody) at a dilution of 1/75(Cytoplasm and Nucleus).

In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with the synthetic peptide and then with 220534(Anti-TDP2 Antibody) at dilution 1/75.



The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 220534(Anti-TDP2 Antibody) at a dilution of 1/75. In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with synthetic peptide and then with D261663(Anti-TDP2 Antibody) at dilution 1/75.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: Mouse small intestines tissue lysate; Primary antibody: 220534(TDP2 Antibody) at dilution 1/600; Secondary antibody: HRP-conjugated Goat anti rabbit IgG at 1/5000 dilution; Exposure time: 90 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010