

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

TEKT1 RABBIT PAB

货号: S210635 产品全名: TEKT1 兔多抗

基因符号

UNIPROT ID: Q969V4 (Gene Accession - BC014599)

背景: This gene product belongs to the tektin family of proteins. Tektins comprise a family of filament-forming proteins that are coassembled with tubulins to form ciliary and flagellar microtubules. This gene is predominantly expressed in the testis and in mouse, tektin 1 mRNA was localized to the spermatocytes and round spermatids in the seminiferous tubules, indicating that it may play a role in spermatogenesis.

抗原: Fusion protein of human TEKT1 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 50-200;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human lymphoma tissue using 210635(TEKT1 Antibody) at a dilution of 1/30(Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human lymphoma tissue is first treated with the fusion protein and then with 210635(Anti-TEKTI Antibody) at dilution 1/30.



The image on the left is immunohistochemistry of paraffinembedded Human cervical cancer tissue using 210635(Anti-TEKTI Antibody) at a dilution of 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with fusion protein and then with D121334(Anti-TEKT1 Antibody) at dilution 1/30.



Gel: 10%SDS-PAGE, Lysate: 40 µg; Lane 1-2: NIH/3T3 cells, Lncap cells; Primary antibody: 210635(TEKT1 Antibody) at dilution 1/500; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 10 minutes



Product Description

Pioneering GTPase and Oncogene Product Development since 2010