

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

TLN1 RABBIT PAB

货号: \$220999 产品全名: TLN1 兔多抗 基因符号 TLN; ILWEQ

UNIPROT ID: Q9Y490 (Gene Accession - NP_006280)

背景: This gene encodes a cytoskeletal protein that is concentrated in areas of cell-substratum and cell-cell contacts. The encoded protein plays a significant role in the assembly of actin filaments and in spreading and migration of various cell types, including fibroblasts and osteoclasts. It codistributes with integrins in the cell surface membrane in order to assist in the attachment of adherent cells to extracellular matrices and of lymphocytes to other cells.

抗原: Synthetic peptide of human TLN1

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Signal Transduction

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 220999(TLNI Antibody) at a dilution of 1/35(Cytoplasm and Cell membrane).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 220999(Anti-TLNI Antibody) at dilution 1/35.



The image on the left is immunohistochemistry of paraffinembedded Human breast cancer tissue using 220999(Anti-TLNI Antibody) at a dilution of 1/35.



In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with synthetic peptide and then with D262338(Anti-TLNI Antibody) at dilution 1/35.

Gel: 6%SDS-PAGE, Lysate: 40 µg; Lane: Huvec cells; Primary antibody: 220999(TLN1 Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010