TNFSF15 RABBIT PAB

货号: S221037
产品全名: TNFSF15 兔多抗
基因符号 TL1; TL1A; VEGI; VEGI192A
UNIPROT ID: O95150 (Gene Accession – NP_005109)
背景: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells.
抗原: Synthetic peptide of human TNFSF15
经过测试的应用: ELISA, WB, IHC
推荐稀释比: IHC: 30-150;WB: 200-1000;ELISA: 1000-2000
种属反应性: Rabbit
克隆性： Rabbit Polyclonal
亚型: Immunogen-specific rabbit IgG
纯化: Antigen affinity purification
种属反应性: Human, Mouse, Rat
成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol
研究领域: Cardiovascular, Immunology
储存和运输: Store at -20°C. Avoid repeated freezing and thawing

Product Description

Pioneering GTPase and Oncogene Product Development since 2010



Immunohistochemistry analysis of paraffin embedded Human liver cancer tissue using 221037(TNFSF15 Antibody) at a dilution of 1/40(Cytoplasm and Cell membrane).



In comparision with the IHC on the left, the same paraffin-embedded Human liver cancer tissue is first treated with the synthetic peptide and then with 221037(Anti-TNFSF15 Antibody) at dilution 1/40.



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using 221037(Anti-TNFSF15 Antibody) at a dilution of 1/40.



In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with synthetic peptide and then with D262387(Anti-TNFSF15 Antibody) at dilution 1/40.



Gel: 12%SDS-PAGE, Lysate: 40 µg;
Lane: Mouse kidney tissue;
Primary antibody: 221037(TNFSF15 Antibody) at dilution 1/600;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 3 minutes