

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## **TREM1 RABBIT PAB**

货号: S217944

产品全名: TREMI 兔多抗 基因符号 CD354; TREM-I

UNIPROT ID: Q9NP99 (Gene Accession - BC017773)

背景: This gene encodes a receptor belonging to the Ig superfamily that is expressed on myeloid cells. This protein amplifies neutrophil and monocyte-mediated inflammatory responses triggered by bacterial and fungal infections by stimulating release of pro-inflammatory chemokines and cytokines, as well as increased surface expression of cell activation markers. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.

抗原: Full length fusion 蛋白 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 2000-5000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Immunology

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 217944(TREM1 Antibody) at a dilution of 1/20(Cytoplasm and Cell membrane).




In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 217944(Anti-TREMI Antibody) at dilution 1/20.



The image on the left is immunohistochemistry of paraffinembedded Human brain tissue using 217944(Anti-TREMI Antibody) at a dilution of 1/20.



In comparision with the IHC on the left, the same paraffin-embedded Human brain tissue is first treated with fusion protein and then with D223411(Anti-TREM1 Antibody) at dilution 1/20.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-2: Hepg2 and 293T cell; Primary antibody: 217944(TREM1 Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 20 seconds



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010