

TRIM27 RABBIT PAB

货号: S217904

产品全名: TRIM27 兔多抗

基因符号: RFP; RNF76

UNIPROT ID: P14373 (Gene Accession - BC013580)

背景: This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein localizes to the nuclear matrix. It interacts with the enhancer of polycomb protein and represses gene transcription. It is also thought to be involved in the differentiation of male germ cells. Fusion of the N-terminus of this protein with the truncated C-terminus of the RET gene product has been shown to result in production of the ret transforming protein.

抗原: Fusion protein of human TRIM27

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 45-300;WB: 500-2000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG


纯化: Antigen affinity purification

种属反应性: Human, Mouse


成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol

研究领域: Epigenetics and Nuclear Signaling


储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Immunohistochemistry analysis of paraffin embedded Human gastric cancer tissue using 217904 (TRIM27 Antibody) at a dilution of 1/65 (Cytoplasm and Nucleus).




In comparison with the IHC on the left, the same paraffin-embedded Human gastric cancer tissue is first treated with the fusion protein and then with 217904 (Anti-TRIM27 Antibody) at dilution 1/65.



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using 217904 (Anti-TRIM27 Antibody) at a dilution of 1/65.



In comparison with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D223342 (Anti-TRIM27 Antibody) at dilution 1/65.



Gel: 8% SDS-PAGE, Lysate: 40 µg;
Lane 1-3: Human testis tissue, 293T and A172 cell lysates;
Primary antibody: 217904 (TRIM27 Antibody) at dilution 1/600;
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;
Exposure time: 10 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
