

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

UCHL3 RABBIT PAB

货号: S217974

产品全名: UCHL3 兔多抗 基因符号 UCH-L3

UNIPROT ID: P15374 (Gene Accession - BC018125)

背景: The protein encoded by this gene is a member of the deubiquitinating enzyme family. Members of this family are proteases that catalyze the removal of ubiquitin from polypeptides and are divided into five classes, depending on the mechanism of catalysis. This protein may hydrolyze the ubiquitinyl-N-epsilon amide bond of ubiquitinated proteins to regenerate ubiquitin for another catalytic cycle. Alternative splicing results in multiple transcript variants that encode different protein isoforms.

抗原: Full length fusion 蛋白 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 500-2000;ELISA: 2000-5000

种属反应性: Rabbit 克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human, Mouse, Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling, Cell Biology 储存和运输: Store at -20°C. Avoid repeated freezing and thawing

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC APPLICATIONS




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human breast cancer tissue using 217974(UCHL3 Antibody) at a dilution of 1/25(Cytoplasm or Nucleus).




In comparision with the IHC on the left, the same paraffin-embedded Human breast cancer tissue is first treated with the fusion protein and then with 217974(Anti-UCHL3 Antibody) at dilution 1/25.



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using 217974(Anti-UCHL3 Antibody) at a dilution of 1/25.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D223472(Anti-UCHL3 Antibody) at dilution 1/25.



Gel: 12%SDS-PAGE, Lysate: 40 µg; Lane 1-4: 293T, K562, Jurkat and 231 cell; Primary antibody: 217974(UCHL3 Antibody) at dilution 1/300; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 30 seconds