

XPNPEP3 ? RABBIT PAB

货号: S213295

产品全名: XPNPEP3 兔多抗

基因符号 APP3; NPHPL1

UNIPROT ID: Q9NQH7 (Gene Accession - NP_071381)

背景: The protein encoded by this gene belongs to the family of X-pro-aminopeptidases that utilize a metal cofactor, and remove the N-terminal amino acid from peptides with a proline residue in the penultimate position. This protein has been shown to localize to the mitochondria of renal cells, and have a role in ciliary function. Mutations in this gene are associated with nephronophthisis-like nephropathy-1. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene, however, expression of some of these isoforms in vivo is not known.

抗原: Synthetic peptide of human XPNPEP3

经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 1000-2000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG



纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg²⁺ and Ca²⁺), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Metabolism

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Immunohistochemistry analysis of paraffin embedded Human lymphoma tissue using 213295 (XPNPEP3 Antibody) at a dilution of 1/40 (Cytoplasm).

In comparison with the IHC on the left, the same paraffin-embedded Human lymphoma tissue is first treated with the synthetic peptide and then with 213295 (Anti-XPNPEP3 Antibody) at dilution 1/40.



Gel: 10% SDS-PAGE, Lysate: 40 µg;

Lane 1-2: MCF7 cells, A431 cells;

Primary antibody: 213295 (XPNPEP3 Antibody) at dilution 1/100;

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution;

Exposure time: 20 seconds