

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

YB1 RABBIT PAB

货号: N225620

产品全名: YBI 兔多抗

基因符号 YBX1; NSEP1; YB1; Nuclease-sensitive element-binding protein 1; CCAAT-binding transcription factor I subunit A; CBF-A; DNA-binding protein B; DBPB; Enhancer factor I subunit A; EFI-A; Y-box transcription factor; Y-boxbinding protein 1; YB-

UNIPROT ID: P67809

背景: This gene encodes a highly conserved cold shock domain protein that has broad nucleic acid binding properties. The encoded protein functions as both a DNA and RNA binding protein and has been implicated in numerous cellular processes including regulation of transcription and translation, pre-mRNA splicing, DNA reparation and mRNA packaging. This protein is also a component of messenger ribonucleoprotein (mRNP) complexes and may have a role in microRNA processing. This protein can be secreted through non-classical pathways and functions as an extracellular mitogen. Aberrant expression of the gene is associated with cancer proliferation in numerous tissues. This gene may be a prognostic marker for poor outcome and drug resistance in certain cancers. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on multiple chromosomes. [provided by RefSeq, Sep 2015]

抗原: A synthesized peptide derived from human YB1

经过测试的应用: WB,IHC-P,ICC/IF,IP,FC

推荐稀释比: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20 FC:

1/50-1/100

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

分子量: Calculated MW: 36 kDa; Observed MW: 50 kDa

亚型: IgG

纯化: Affinity Chromatography

种属反应性: Human, Mouse and Rat

成分: PBS (without Mg2+ and Ca2+), pH 7.3 containing 50% glycerol,

0.5% BSA and 0.02% sodium azide

研究领域: Tags & Cell Markers

储存和运输: Store at -20°C. Avoid repeated freezing and thawing



Product Description

Pioneering GTPase and Oncogene Product Development since 2010





Immunohistochemistry analysis of paraffin-embedded Human kidney using YB1 antibody.High- V

kidney using YB1 antibody.High- Western blot analysis of YB1 in pressure and temperature Sodium HeLa lysates using YB1 antibody. Citrate pH 6.0 was used for antigen retrieval.



Immunofluorescence analysis of YBI in MCF-7 using YBI antibody.