

货号: S218773

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010


## **ZBED1 RABBIT PAB**

产品全名: ZBED1 兔多抗 基因符号 ALTE; DREF; TRAMP; hDREF UNIPROT ID: 096006 (Gene Accession - BC015030) 背景: This gene is located in the pseudoautosomal region 1 (PAR1) of X and Y chromosomes. It was earlier identified as a gene with similarity to Ac transposable elements, however, was found not to have transposase activity. Later studies show that this gene product is localized in the nucleus and functions as a transcription factor. It binds to DNA elements found in the promoter regions of several genes related to cell proliferation, such as histone HI, hence may have a role in regulating genes related to cell proliferation. Alternatively spliced transcript variants with different 5' untranslated region have been found for this gene. 抗原: Fusion protein of human ZBED1 经过测试的应用: ELISA, WB, IHC 推荐稀释比: IHC: 50-300;WB: 200-1000;ELISA: 2000-5000 种属反应性: Rabbit 克隆性: Rabbit Polyclonal 亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification 种属反应性: Human 成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol 研究领域: Epigenetics and Nuclear Signaling 储存和运输: Store at -20°C. Avoid repeated freezing and thawing




## **Product Description**


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human thyroid cancer tissue using 218773(ZBED1 Antibody) at a dilution of 1/50(Nucleus).



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with the fusion protein and then with 218773(Anti-ZBEDI Antibody) at dilution 1/50.



The image on the left is immunohistochemistry of paraffinembedded Human cervical cancer tissue using 218773(Anti-ZBED1 Antibody) at a dilution of 1/50.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane: A549 cell lysate; Primary antibody: 218773(ZBEDI Antibody) at dilution 1/350; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 1 minute

In comparision with the IHC on the left, the same paraffin-embedded Human cervical cancer tissue is first treated with fusion protein and then with D225146(Anti-ZBEDI Antibody) at dilution 1/50.



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010