

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

ZNF8 RABBIT PAB

货号: S218256

产品全名: ZNF8 兔多抗 基因符号 HF.18; Zfp128

UNIPROT ID: P17098 (Gene Accession - BC039323)

背景: ZNF8 (zinc finger protein 8) is a 575 amino acid protein that belongs to the KrIppel C2H2-type zinc-finger protein family and contains seven C2H2-type zinc fingers and a KRAB domain. Localizing to the nucleus and ubiquitously expressed, ZNF8 is thought to play a role in transcriptional regulation and is encoded by a gene that maps to human chromosome 19q13.43.

抗原: Fusion protein of human ZNF8 经过测试的应用: ELISA, WB, IHC

推荐稀释比: IHC: 25-100;WB: 200-1000;ELISA: 5000-10000

种属反应性: Rabbit

克隆性: Rabbit Polyclonal

亚型: Immunogen-specific rabbit IgG 纯化: Antigen affinity purification

种属反应性: Human

成分: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.05% Sodium Azide and 40% glycerol


研究领域: Epigenetics and Nuclear Signaling

储存和运输: Store at -20°C. Avoid repeated freezing and thawing




Product Description


Pioneering GTPase and Oncogene Product Development since 2010




Immunohistochemistry analysis of paraffin embedded Human esophagus cancer tissue using 218256(ZNF8 Antibody) at a dilution of 1/30(Nucleus or Cytoplasm).




In comparision with the IHC on the left, the same paraffin-embedded Human esophagus cancer tissue is first treated with the fusion protein and then with 218256(Anti-ZNF8 Antibody) at dilution 1/30.



The image on the left is immunohistochemistry of paraffinembedded Human thyroid cancer tissue using 218256(Anti-ZNF8 Antibody) at a dilution of 1/30.



In comparision with the IHC on the left, the same paraffin-embedded Human thyroid cancer tissue is first treated with fusion protein and then with D224045(Anti-ZNF8 Antibody) at dilution 1/30.



Gel: 8%SDS-PAGE, Lysate: 40 µg; Lane 1-3: SKOV3, A549 and HEPG2 cell lysates; Primary antibody: 218256(ZNF8 Antibody) at dilution 1/200; Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution; Exposure time: 5 seconds



Product Description

Pioneering GTPase and Oncogene Product Development since 2010