

HUMAN CCR3 FULL LENGTH PROTEIN

货号: 11056

产品全名: 人 CCR3 Full Length 蛋白

规格: 10/50/100 µg

基因符号 CC-CKR-3; C C CKR3; CD193; CKR 3; CKR3; CMKBR3

目标蛋白: CCR3

UNIPROT ID: P51677

描述: 人 CCR3 full length protein-synthetic nanodisc

背景: The protein is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced transcript variants have been described.

物种/宿主: HEK293

分子量: The human full length CCR3 protein has a MW of 41.0 kDa

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

ELISA assay to evaluate CCR3-Nanodisc
0.2µg Human CCR3-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CCR3-Nanodisc (0.2 µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR3-Nanodisc is 2.3ng/ml.



Figure2. Human CCR3-Nanodisc, Flag Tag on SDS-PAGE