

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN CD37 FULL LENGTH PROTEIN

货号: 11023 产品全名: 人 CD37 Full Length 蛋白 规格: 10/50/100 μg 基因符号 GP52-40; TSPAN26 目标蛋白: CD37

UNIPROT ID: P11049

描述:人CD37 full length protein-synthetic nanodisc

背景: The protein is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins. It may play a role in T-cell-B-cell interactions. Alternate splicing results in multiple transcript variants encoding different isoforms.

物种/宿主: HEK293

分子量: The human full length CD37 protein has a MW of 31.7 kDa

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Figure1. Elisa plates were pre-coated with Flag Tag CD37-Nanodisc (0.2 µg/per well). Serial diluted anti-CD37 monoclonal antibody (28055) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CD37 monoclonal antibody binding with CD37-Nanodisc is 23.06ng/ml.




Figure2. Human CD37-Nanodisc, Flag Tag on SDS-PAGE




Figure3. WB analysis of Human CD37-Nanodisc with anti-CD37 monoclonal antibody (28055), followed by Goat Anti-Human IgG HRP at 1/5000 dilution