

## EYA2 MAB

产品名称: EYA2 小鼠单抗

货号: 26096

基因符号: EYA2

描述: EYA2 小鼠单抗

背景: EYA2 may be involved in development of the eye. It promotes efficient DNA repair by dephosphorylating H2AX, promoting the recruitment of DNA repair complexes containing MDC1. Its function as histone phosphatase probably explains its role in transcription regulation during organogenesis. EYA2 coactivates SIX1 and seems to coactivate SIX2, SIX4 and SIX5. It seems to be involved in myogenesis together with SIX1 and DACH2.

免疫原: 人 EYA2 蛋白

经过测试的应用: ELISA, WB, IHC

推荐稀释比:

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IHC: 1:50-1:100

**Concentration:** 0.8 mg/ml

种属反应性: Mouse

形式: Liquid

克隆性: Monoclonal

亚型: IgG

纯化: Purified from ascites


成分: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 50% glycerol  
Species

种属反应性: EYA2 antibody recognizes EYA2 of vertebrates.

储存条件: Store at -20°C. Avoid repeated freezing and thawing.

## 蛋白质印迹:


EYA2 protein



WB: Anti-EYA2 mAb

Western blot analysis of recombinant full length EYA2 protein. Purified His-tagged EYA2 protein(货号:10195) was blotted with anti EYA2 monoclonal antibody (Cat # 26096).

## Immunohistochemistry:



Immunohistochemical analysis of paraffin-embedded thyroid cancer tissue with anti EYA2 monoclonal antibody (货号:26096). Tissue samples were fixed with formaldehyde and blocked with 1% serum for 15 min at 37 °C. Antigen retrieval was by heat mediation in citrate buffer (pH6). Samples were then incubated with primary antibody (1:50) overnight at 4°C. A HRP-conjugated Goat anti-mouse IgG (dilution 1:50) was used as secondary antibody.