

HUMAN CCL16 PROTEIN

货号: 12100

产品全名: 人 CCL16 蛋白

规格: 10/50/100 µg

基因符号 C-C Motif Chemokine 16;Chemokine CC-4;HCC-4;Chemokine LEC;IL-10-Inducible Chemokine;LCC-1;Liver-Expressed

Chemokine;Lymphocyte and Monocyte

Chemoattractant;LMC;Monotactin-1;MTN-1;NCC-4;Small-Inducible Cytokine A16;CCL16;ILINCK;NCC4;SCYA16

目标蛋白: CCL16

UNIPROT ID: O15467

描述: Recombinant Human C-C Motif Chemokine 16 is produced by our E.coli expression system and the target gene encoding Gln24-Gln120 is expressed.

背景: CCL16 is a member of CC chemokine family. CCL16 cDNA encodes a 120 amino acid peptide along with a 23 amino acids signal peptide that is cleaved to generate 97 amino acid protein. CCL16 is distantly related to other CC chemokines, showing less than 30% sequence identity. CCL16 elicits its effects on cells by interacting with cell surface chemokine receptors such as CCR1, CCR2, CCR5 and CCR8. Recombinant CCL16 has been shown to chemoattract human monocytes and THP1 cells but not resting lymphocytes nor neutrophils. CCL16 has potent myelosuppressive activity, suppresses proliferation of myeloid progenitor cells. CCL16 induces a calcium flux in THP1 cells that can be desensitized by prior exposure to RANTES, suggesting that CCL16 and RANTES share the same receptor in THP1 cells.

物种/宿主: E.coli

分子量: 11 KDa

分子特征: Not available

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010
