

HUMAN CD32A (H167) PROTEIN, HFC TAG

货号: 11469

产品全名: 人 CD32A (H167) 蛋白

规格: 10/50/100 µg

基因符号 CD32;CD32A;CDw32;FCG2;FcGR;FCGR2;FCGR2A1;IGFR2

目标蛋白: CD32α

UNIPROT ID: P12318

描述: Recombinant Human CD32A(167H) with C-terminal human Fc tag

背景: This gene encodes one member of a family of immunoglobulin Fc receptor genes found on the surface of many immune response cells. The protein encoded by this gene is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2008]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 46.4 kDa after removal of the signal peptide. The apparent molecular mass of CD32A(167H)-hFc is approximately 55-70 kDa due to glycosylation.

分子特征: CD32α(H167)(Ala36-Ile218) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CD32 α (H167) Protein, hFc Tag on SDS-PAGE under reducing condition.