

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN CLDN18.2 FULL LENGTH PROTEIN

目录: 12238

产品名称: Human CLDN18.2 Full Length Protein

规格: 10 μg, 50 μg and 100 μg

基因符号: Claudin 18.2

Target: CLDN18.2 UNIPROT ID: P56856

描述: Human Full length Claudin 18.2 Protein-VLP

背景: The protein encodes a member of the claudin family. Claudins are integral membrane proteins and components of tight junction strands. Tight junction strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. This gene is upregulated in patients with ulcerative colitis and highly overexpressed in infiltrating ductal adenocarcinomas. PKC/MAPK/AP-1 (protein kinase C/mitogen-activated protein kinase/activator protein-1) dependent pathway regulates the expression of this gene in gastric cells. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jun 2010]

Species/Host: HEK293

Molecular Weight: The Human CLDN18.2 Protein has a predicted MW of 29 kDa.Due to PTM, the actual MW on SDS-PAGE gel is around 50kDa.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Figure I. ELISA plates were pre-coated with 0.5 ug/per well purified human CLDN 18.2 VLP. Serial diluted CLDN 18.2 monoclonal antibody (Zolbetuximab biosililar; IMAB 362) solutions were added, washed, and incubated with secondary antibody before ELISA reading. From above data, the EC50 for IMAB 362 binding with CLDN 18.2 VLP is 15.37 ng/ml.