

HUMAN CXCR3 PROTEIN, MFC TAG

货号: 11585

产品全名: 人 CXCR3 蛋白

规格: 10/50/100 µg

基因符号 CXC-R3;CXCR-3;CKR-L2;IP-10 receptor;CD183;GPR9

目标蛋白: CXCR3

UNIPROT ID: P49682

描述: Recombinant human CXCR3 protein with C-terminal mouse Fc tag

背景: This gene encodes a G protein-coupled receptor with selectivity for three chemokines, termed CXCL9/Mig (monokine induced by interferon- γ), CXCL10/IP10 (interferon- γ -inducible 10 kDa protein) and CXCL11/I-TAC (interferon-inducible T cell α -chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic, most notably integrin activation, cytoskeletal changes and chemotactic migration. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine, CXCL4/PF4 (PMID:12782716). [provided by RefSeq, Jun 2011]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 32.1 kDa after removal of the signal peptide. The apparent molecular mass of CXCR3-mFc is approximately 25–55 kDa due to glycosylation.

分子特征: CXCR3(Met1-Arg53) mFc(Pro99-Lys330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Figure 1. Human CXCR3 Protein, mFc Tag on SDS-PAGE under reducing condition.

Human CXCR3,mFc Tagged protein ELISA

0.2 µg of Human CXCR3, mFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CXCR3 Protein, mFc Tag(11585) can bind Anti-CXCR3 antibody(DM208), Rabbit mAb in a linear range of 0.13–3.20 ng/mL.