

## HUMAN FLT3 PROTEIN, HFC-HIS TAG

货号: 11135

产品全名: 人 FLT3 蛋白

规格: 10/50/100  $\mu$ g

基因符号 Flt-3;Flk-2;STK-1;CD135;FLK2;FLT-3

目标蛋白: FLT3

**UNIPROT ID:** P36888

**描述:** Recombinant human FLT3 protein with C-terminal human Fc and 6xHis tag

**背景:** Flt-3(Receptor-type tyrosine-protein kinase FLT3) is also known as FLK-2(Fetal liver kinase-2), STK-1(Stem cell tyrosine kinase 1), CD135. FLT3 is a cytokine receptor which belongs to the receptor tyrosine kinase class III. Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Mutations that cause constitutive kinase activity promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways.

**物种/宿主:** HEK293

**分子量:** The protein has a predicted molecular mass of 125-130 kDa after removal of the signal peptide.

**分子特征:** FLT3(Asn27-Ser543) hFc(Glu99-Ala330) 6xHis tag

**纯化:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization.

**储存和运输:** Store at  $-20^{\circ}\text{C}$  to  $-80^{\circ}\text{C}$  for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at  $-80^{\circ}\text{C}$  (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human FLT3, hFc-His Tag on SDS-PAGE under reducing condition.

### Human FLT3, hFc-His Tagged protein ELISA

0.2  $\mu$ g of Human FLT3LG, mFc-His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human FLT3LG, mFc-His tagged protein [11161] can bind Human FLT3, hFc-His tagged protein (11135) in a linear range of 0.128-10.02 ng/ml.



# Product Description

Pioneering GTPase and Oncogene Product Development since 2010

---