

HUMAN GDF-5 PROTEIN

货号: 12059

产品全名: 人 GDF-5 蛋白

规格: 10/50/100 µg

基因符号 Growth/differentiation factor 5;GDF-5;Bone morphogenetic protein 14;BMP-14;Cartilage-derived morphogenetic protein 1;CDMP-1;Lipopolysaccharide-associated protein 4;LAP-4;LPS-associated protein 4;Radotermis;CDMP1

目标蛋白: GDF-5

UNIPROT ID: P43026

描述: Recombinant Human Growth/Differentiation Factor 5 is produced by our E.coli expression system and the target gene encoding Ala382-Arg501 is expressed.

背景: Growth Differentiation Factor 5(GDF-5, BMP-14) is a member of the BMP family of TGFβ superfamily proteins. Human GDF-5, -6, and -7 are a defined subgroup of the BMP family. GDF-5 is synthesized as a homodimeric precursor protein consisting of a 354 amino acid (aa) Nterminal proregion and a 120 aa C-terminal mature peptide. Mature human GDF-5 shares 99% aa sequence identity with both mature mouse and rat GDF-5. GDF-5 signaling is mediated by formation of a heterodimeric complex consisting of a type I (BMPRII) and a type II (BMPRI or Activin RII) serine/threonine kinase receptor which results in the phosphorylation and activation of cytosolic Smad proteins (Smad1, 5, and 8). GDF-5 is involved in multiple developmental processes including limb generation, cartilage development, joint formation, bone morphogenesis, cell survival, and neuritogenesis. Inhibition of GDF-5 expression or alteration of its signaling can facilitate the development of osteoarthritis.

物种/宿主: E.coli

分子量: 13.7 KDa

分子特征: Not available

纯化: Greater than 95% as determined by reducing SDS-PAGE.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Greater than 95% as determined by reducing SDS-PAGE.