

HUMAN GP1BB PROTEIN, HFC TAG

货号: 11428

产品全名: 人 GP1BB 蛋白

规格: 10/50/100 µg

基因符号 BDPLT1;BS;CD42C;GP1BB;GP1bbeta

目标蛋白: GP1BB

UNIPROT ID: P13224

描述: Recombinant Human GP1BB with C-terminal human Fc tag

背景: Platelet glycoprotein Ib (GP1b) is a heterodimeric transmembrane protein consisting of a disulfide-linked 140 kD alpha chain and 22 kD beta chain. It is part of the GP1b-V-IX system that constitutes the receptor for von Willebrand factor (VWF), and mediates platelet adhesion in the arterial circulation. GP1b alpha chain provides the VWF binding site, and GP1b beta contributes to surface expression of the receptor and participates in transmembrane signaling through phosphorylation of its intracellular domain. Mutations in the GP1b beta subunit have been associated with Bernard-Soulier syndrome, velocardiofacial syndrome and giant platelet disorder. The 206 amino acid precursor of GP1b beta is synthesized from a 1.0 kb mRNA expressed in platelets and megakaryocytes. A 411 amino acid protein arising from a longer, unspliced transcript in endothelial cells has been described; however, the authenticity of this product has been questioned. Yet another less abundant GP1b beta mRNA species of 3.5 kb, expressed in nonhematopoietic tissues such as endothelium, brain and heart, was shown to result from inefficient usage of a non-consensus polyA signal in the neighboring upstream gene (SEPT5, septin 5). In the absence of polyadenylation from its own imperfect site, the SEPT5 gene produces read-through transcripts that use the consensus polyA signal of this gene.

[provided by RefSeq, Dec 2010]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 39.0 kDa after removal of the signal peptide. The apparent molecular mass of GP1BB-hFc is approximately 35-55 kDa due to glycosylation.

分子特征: GP1BB(Pro27-Cys147) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human GPIBB Protein, hFc Tag on SDS-PAGE under reducing condition.