

HUMAN IL19 PROTEIN, HFC TAG

货号: 11907

产品全名: 人 IL19 蛋白

规格: 10/50/100 µg

基因符号 IL-10C;MDA1;NG.1;ZMDA1

目标蛋白: IL19

UNIPROT ID: Q9UHD0

描述: Recombinant Human IL19 Protein with C-terminal human Fc tag

背景: The protein encoded by this gene is a cytokine that belongs to the IL10 cytokine subfamily. This cytokine is found to be preferentially expressed in monocytes. It can bind the IL20 receptor complex and lead to the activation of the signal transducer and activator of transcription 3 (STAT3). A similar cytokine in mouse is reported to up-regulate the expression of IL6 and TNF-alpha and induce apoptosis, which suggests a role of this cytokine in inflammatory responses. Alternatively spliced transcript variants encoding the distinct isoforms have been described. [provided by RefSeq, Jul 2008]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 43.9 kDa after removal of the signal peptide. The apparent molecular mass of IL19-hFc is approximately 35-70 kDa due to glycosylation.

分子特征: IL19(Leu25-Ala177) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Figure 1. Human IL19 Protein, hFc Tag on SDS-PAGE under reducing condition.