

HUMAN IL22 PROTEIN, HFC TAG

货号: 11447

产品全名: 人 IL22 蛋白

规格: 10/50/100 µg

基因符号 IL-22;L-TIF;CYTO18

目标蛋白: IL22

UNIPROT ID: Q9GZX6

描述: Recombinant Human IL22 with C-terminal human Fc tag

背景: This gene is a member of the IL10 family of cytokines that mediate cellular inflammatory responses. The encoded protein functions in antimicrobial defense at mucosal surfaces and in tissue repair. This protein also has pro-inflammatory properties and plays a role in the pathogenesis of several intestinal diseases. [provided by RefSeq, Jul 2018]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 33.9 kDa after removal of the signal peptide. The apparent molecular mass of IL22-hFc is approximately 35-55 kDa due to glycosylation.

分子特征: IL22(Ala34-Ile179) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Figure 1. Human IL22 Protein, hFc Tag on SDS-PAGE under reducing condition.

Human IL22, hFc Tagged protein ELISA

0.2 μ g of Human IL22, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human IL22 Protein, hFc Tag (11447) can bind Anti-IL22 Neutralizing antibody 28112 in a linear range of 2.44-625.00 ng/mL.