

HUMAN LOX-1 PROTEIN, HFC TAG

货号: 11968

产品全名: 人 LOX-1 蛋白

规格: 10/50/100 µg

基因符号 CLEC8A;LOX1;LOXIN;SCARE1;SLOX1;OLRI

目标蛋白: LOX-1

UNIPROT ID: P78380

描述: Recombinant Human LOX-1 Protein with N-terminal human Fc tag

背景: This gene encodes a low density lipoprotein receptor that belongs to the C-type lectin superfamily. This gene is regulated through the cyclic AMP signaling pathway. The encoded protein binds, internalizes and degrades oxidized low-density lipoprotein. This protein may be involved in the regulation of Fas-induced apoptosis. This protein may play a role as a scavenger receptor. Mutations of this gene have been associated with atherosclerosis, risk of myocardial infarction, and may modify the risk of Alzheimer's disease. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Feb 2010]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 50.9 kDa after removal of the signal peptide. The apparent molecular mass of hFc-LOX-1 is approximately 55-70 kDa due to glycosylation.

分子特征: hFc(Glu99-Ala330) LOX-1(Met58-Gln273)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Figure 1. Human LOX-1 Protein, hFc Tag on SDS-PAGE under reducing condition.