

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN MICB PROTEIN, HIS TAG

货号: 11318

产品全名: 人 MICB 蛋白 规格: 10/50/100 µg

基因符号 MIC-B;PERBII.2

目标蛋白: MICB

UNIPROT ID: Q29980

描述: Recombinant Human MICB Protein with C-terminal 6xHis tag

背景: This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells, CD8 alphabeta T cells, and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules, however, it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 33.5 kDa after removal of the signal peptide. The apparent molecular mass of MICB-His is approximately 35-55 kDa due to glycosylation.

分子特征: MICB(Ala23-Asp309) 6×His tag

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. Human MICB Protein, His Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human MICB protein, His Tag (11318) can bind Anti-MICB Rabbit mAb in a linear range of 0.128-16 ng/mL.




Figure 3. Flow cytometry analysis with 15 µg/ml Human MICB Protein, His Tag (11318) on Expi293 cells transfected with Human NKG2D protein and Human DAP10 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).