

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN MST1R PROTEIN, HFC TAG

货号: 11602

产品全名: 人 MSTIR 蛋白

规格: 10/50/100 µg

基因符号 CD136;CDw136;NPCA3;PTK8;RON

目标蛋白: MSTIR

UNIPROT ID: Q04912

描述: Recombinant Human MSTIR Protein with C-terminal Human Fc tag

背景: This gene encodes a cell surface receptor for macrophage-stimulating protein (MSP) with tyrosine kinase activity. The mature form of this protein is a heterodimer of disulfide-linked alpha and beta subunits, generated by proteolytic cleavage of a single-chain precursor. The beta subunit undergoes tyrosine phosphorylation upon stimulation by MSP. This protein is expressed on the ciliated epithelia of the mucociliary transport apparatus of the lung, and together with MSP, thought to be involved in host defense. Alternative splicing generates multiple transcript variants encoding different isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Jan 2016]

物种/宿主: HEK293

分子量: The protein has a predicted molecular mass of 125.9 kDa after removal of the signal peptide. The apparent molecular mass of MSTIR-hFc is approximately 130-250 kDa due to glycosylation.

分子特征: MSTIR(Glu25-Thr957) hFc(Glu99-Ala330)

纯化: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

储存和运输: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



Product Description

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. Human MST1R Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human MST1R Protein, hFc Tag (11602) can bind Anti-MST1R Neutralizing antibody 28116 in a linear range of 0.73-46.88 ng/mL.