HUMAN TGFB3 PROTEIN, HFC TAG

： 11718

： TGFB3

： 10/50/100 μg

ARVD;ARVD1;LDS5;RNHF;TGF-beta3

： TGFB3

UNIPROT ID: P10600

： Recombinant human TGFB3 protein with C-terminal human Fc tag

： This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. This protein is involved in embryogenesis and cell differentiation, and may play a role in wound healing. Mutations in this gene are a cause of aortic aneurysms and dissections, as well as familial arrhythmogenic right ventricular dysplasia 1. [provided by RefSeq, Aug 2016]

／ ： HEK293

： The protein has a predicted molecular mass of 56 kDa after removal of the signal peptide.

： TGFB3 (Leu24-Arg300) hFc (Glu99-Ala330)

： The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

： Store at -20ºC to -80ºC for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80ºC (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.